

Depletion of human microRNA miR-125b reveals that it is critical for the proliferation of differentiated cells but not for the down-regulation of putative targets during differentiation

Yong Sun Lee¹, Hak Kyun Kim^{1,2}, Sangmi Chung³, Kwang-Soo Kim³, and Anindya Dutta^{1,4}

1. Department of Biochemistry & Molecular Genetics, University of Virginia, Charlottesville, VA 22908, United States
2. Institute of Molecular Biology and Genetics, School of Biological Sciences, Seoul National University, Seoul 151-742, Korea.
3. Molecular Neurobiology Laboratory, McLean Hospital, Harvard Medical School, Belmont, MA 02478, United States
4. Corresponding author; 434-924-1227 (phone), 434-924-5069 (fax), ad8q@virginia.edu (e-mail)

Short running title: miR-125b during differentiation

Keywords: microRNA, differentiation, miR-125b, PC-3 cell line, proliferation, lin-28, transcriptional regulation

Summary

MicroRNAs are small non-coding RNAs that regulate target gene expression post-transcriptionally through base pairing with the target messenger RNA. Functional characterization of microRNAs awaits robust experimental methods to knock-down a microRNA as well as to assay its function *in vivo*. In addition to the recently developed method to sequester microRNA with 2'-O-methyl antisense oligonucleotide, we report that siRNA against the loop region of a microRNA precursor can be used to deplete the microRNA. The depletion of miR-125b by this method had a profound effect on the proliferation of adult differentiated cancer cells and this proliferation defect was rescued by co-transfected mature microRNA. This technique has unique advantages over the 2'-O-methyl antisense oligonucleotide and can be used to determine microRNA function, assay microRNAs *in vivo* and identify the contribution of a predicted microRNA precursor to the pool of mature microRNA in a given cell. miR-125b and *let-7* microRNAs are induced whereas their putative targets, *lin-28* and *lin-41*, are decreased during *in vitro* differentiation of Tera-2 or embryonic stem cells. Experimental increase or decrease of microRNA concentrations did not, however, affect the levels of the targets a finding that is explained by the fact that the downregulation of the targets appear to be mostly at the transcriptional level in these *in vitro* differentiation systems. Collectively these results reveal the importance of microRNA depletion strategies for directly determining microRNA function *in vivo*.

Introduction

MicroRNAs (miRNAs) are a class of small non-coding RNAs that are processed by Dicer from precursors with a characteristic hairpin secondary structure (1). It is now known that there are hundreds of evolutionarily conserved miRNAs in plants and animals, but their functions are not well understood (2-16). Current knowledge of miRNA function is mainly based on the result from the founding members of miRNA, *lin-4* (2) and *let-7* (17) in *C. elegans*. Both are heterochronic gene whose mutation results in developmental defect (reviewed in (18)). Up-regulation of *lin-4* during the first larval stage (L1) triggers the subsequent developmental stages by translational repression of *lin-14* and *lin-28* both of which contain sites of partial complementary to *lin-4* in the 3' untranslated region (19,20). Likewise, *let-7* is induced during development and translationally repress *lin-41* through partially complementary base pairing between *let-7* and target sequences in the 3' untranslated region of *lin-41* (17,21-23).

The study of a miRNA function has been hampered by the lack of an efficient method to deplete a miRNA, especially in mammals where genetic experiments are hard to perform. Kawasaki and Taira (24) reported a knock-down of a miRNA by a siRNA against the loop region of the miRNA precursor, but the paper was retracted (25). Recently, 2'-O-methyl antisense oligonucleotide against a miRNA was reported to specifically knock-down the miRNA (26,27). Despite the successful knock-down of a miRNA *in vitro* and *in vivo*, this method has several limitations. First, a direct measurement of the depletion of a miRNA is difficult, because 2'-O-methyl antisense oligonucleotide binds to the miRNA and sequesters it from its target, rather than induce its degradation. In addition, even, a slight contamination of 2'-O-methyl antisense

oligonucleotide in an RNA preparation interferes with assays to measure the miRNA level, such as Northern blot, primer extension assay, or RNase protection assay (YSL and AD, unpublished). Therefore, the only possible way to confirm the decrease of a miRNA is to employ a very indirect and cumbersome assay: measuring the level of expression of a reporter gene containing the target sequence of the miRNA. In addition, although rescuing the depletion effect by adding back miRNA is the most stringent control against nonspecific effects of miRNA depletion, the knockdown phenotype cannot be rescued in the presence of the 2'-O-methyl antisense oligonucleotide.

In addition to 2'-O-methyl antisense oligonucleotides to miRNAs, we now show that siRNA against the loop region of a pre-miRNA can be used to specifically downregulate miRNAs. The unique advantages of this method over 2'-O-methyl antisense oligonucleotide will be discussed. Using both these methods, we show that miR-125b, putative homolog of *lin-4* in *C. elegans* (9), is critical for the proliferation of differentiated human cell lines, providing a new example of the biological role of a mammalian miRNA. In addition, we could restore miRNA function in the miRNA-depleted cells by introduction of single-stranded RNA representing the miRNA sequence.

Although the exact regulatory mechanism of mammalian *lin-28* and *lin-41* in *in vitro* differentiation systems or the role of the miRNAs in this regulation has not yet been described, a translational control of mammalian Lin-28 has been hypothesized similar to that seen in *C. elegans* (28-30). The hypothesis is based on the following observations: 1) the evolutionary conservation of *lin-4* and *let-7* miRNAs and their targets *lin-28* and *lin-41*, 2) the presence of putative target sequences with imperfect complementarity to miR-125b and *let-7* within the 3' UTR of *lin-28* and *lin-41* and 3) induction of miR-125b

and *let-7* and the attendant decrease of Lin-28 protein during differentiation. This assumption was supported by an experiment with artificial luciferase construct fused to the *let-7* target sequence that is naturally found in the 3' UTR of mammalian *lin-28* mRNAs (30). Using the methods developed above to decrease or increase miR-125b or *let-7* we, however, find that the decrease of *lin-28* and *lin-41* during differentiation appears independent of the levels of these miRNAs, demonstrating the importance of miRNA depletion strategies to identify real targets of miRNAs *in vivo*. The negative result is probably explained by the fact that the decrease of *lin-28* and *lin-41* is mainly at the transcriptional stage, suggesting a variation in the paradigm of *lin-28* and *lin-41* regulation during differentiation of mammalian cells *in vitro*.

Experimental Procedures

RNA isolation, Northern blotting, and primer extension assay

Total RNAs from various cell lines were isolated with Trizol reagent (Life Technologies) according to the manufacturer's instructions. Total RNAs from human cervix, prostate, and brain were purchased from Ambion. Multiple tissue Northern blot (human MTN blot II) was purchased from Clontech.

Primer extension assay (31) was performed with Superscript II reverse transcriptase (Invitrogen Life Technologies) as manufacturer's instructions with minor modifications. 20 μ l of extension reaction contained 0.1 pmol of the γ -³²P-labeled primer and 10 μ g of total RNA. In the case of 5S rRNA extension reaction, 1 μ g of total RNA was used. Extension reaction at 50°C for 50 minutes was quenched by addition of 20 μ l of sequencing gel loading buffer (32). Extension products were resolved from the primer by electrophoresis in an 18% polyacrylamide gel with 7M urea.

The sequences of primers are as following;

let-7a-3-ext: 5'GCCCCAAACTATAACAACCTACTAC3'

miR-125a-ext: 5'TCACAGGTAAAGGGTCTCA3'

miR-125b-ext: 5'CATCACAAGTTAGGGTCTCA3'

5S rRNA-ext: 5'GATCGGGCGCGTTCAGGGTGGTAT3'

miR-16-ext: 5'ACTACGCCAATATTTACGTGCT3'

miR-15a-ext: 5'AATCCACAAACCATTATGTGCT3'

siRNAs, 2'-O-methyl oligonucleotides, miRNA, and transfections

siRNA duplexes and 2'-O-methyl oligonucleotides were synthesized by Dharmacon. Transfection was performed with oligofectamine reagent (Invitrogen Life Technologies) as manufacturer's instructions. Unless indicated, 960 nM of siRNA duplexes designed against loop regions of miRNAs was used for the transfection into Tera-2 or PC-3 cell lines (ATCC No.; CRL-1435). In the case of 2'-O-methyl antisense oligonucleotides against miR-125b or GL2, 160 nM of oligonucleotides was transfected into PC3 or HeLa cells.

The target sequences for siRNA duplexes are as following;

siRNA against miR-125b-1 (si125b-1): 5'AACCGUUUAAAUCCACGGGUU3'

siRNA against miR-125b-2 (si125b-2): 5'AAGGUAUUUUAGUAACAUCAC3'

siRNA against miR-16-1 (si16-1): 5'AAGAUUCUAAAAUUAUCUCCA3'

siRNA against miR-16-2 (si16-2): 5'GGCGUAGUGAAAUAUAUAU3'

siRNA against miR-15a (si15a): 5'GUGGAUUUUGAAAAGGUGC3'

RNA oligonucleotides of mature let-7a-3 and miR-125b sequences were synthesized by Dharmacon. Transfection was performed as described above, but at the final concentration of 2 μ M. During long-term experiment with multiple transfections, we performed first three transfections everyday and the transfections from the fourth and later on every other day.

Cell proliferation assay

Cell growth was measured with CellTiter 96 non-radioactive cell proliferation assay kit (Promega).

Cell culture, retinoic acid treatment, and stem cell differentiation

Tera-2 cells (ATCC No.; HTB-106) were treated with 5 μ M of all-trans-retinoic acid (Sigma) on gelatin-coated tissue culture dish (33). Culture medium with retinoic acid was refreshed every other day. The cells were cultured in the presence of retinoic acid for 20 days, with subsequent cultivation for additional 40 days without retinoic acid.

Neural induction from mouse ES cell line D3 (ATCC No.; CRL-11632) was maintained and differentiated using the 5-stage procedure as described (34). Briefly, undifferentiated ES cells were cultured on gelatin-coated dishes in supplemented DMEM (Life Technologies, Rockville, MD), and differentiated into EB on nonadherent bacterial dishes. The EBs were then plated onto adhesive tissue culture surface, and neuronal precursor (NP) cells were selected in serum-free medium. After 6 to 10 days of selection, cells were trypsinized and nestin⁺ NP cells were expanded in N2 medium supplemented with 1 μ g/ml laminin (Sigma) and 10 ng/ml bFGF (R & D Systems). After expansion for four days, bFGF was removed to induce differentiation to neuronal phenotypes.

Nuclear run-on assay

ORFs of *lin-28* and *lin-41* were PCR-amplified from Tera-2 cDNA. 0.5 μ g of each fragment was slot blotted onto positively charged Nylon membrane (Nytran from Schleicher & Schuell).

Nucleus isolation and run-on transcription reaction were performed as described (35) with modifications. After the transcription reaction, Trizol reagent (Life Technologies) was added, and RNA probe was prepared according to the manufacturer's instructions for Trizol reagent. Membranes were probed with ³²P-labeled run-on RNA for

16 h at 68°C in hybridization buffer containing 0.25 M Na₂HPO₄ (pH 7.2), 1 mM EDTA and 7% (w/v) SDS. Membranes were washed twice in 2x SSC at room temperature for 15 min each, followed by a wash in 0.2x SSC and 1% (w/v) SDS at 65°C for 20 min. The blot was exposed to phosphorimager Storm 860 (Molecular Dynamics) and quantitated.

Recombinant Lin-28 and antiserum

cDNA containing full-length human Lin-28 protein (209 amino acids) was cloned into pET-23a(+) (Novagen). Recombinant Lin-28 with hexahistidine tag was expressed in BL21(DE3), purified through nickel-charged chelating sepharose column (Amersham Biosciences), and used for raising antiserum in rabbit. Antibody was proven specific by the following criteria. It recognized the recombinant Lin-28 and a single protein band of expected size of Lin-28 in Tera-2 cell extract, whereas preimmune serum did not detect protein of appropriate size (data not shown). In addition, it failed to detect protein in the extract of PC-3 cells where *lin-28* mRNA was not detectable.

Results

Expression of *let-7* and miR-125b in differentiated cells

Primer extension assays were employed to quantitate miR-125a and miR-125b, miRNAs that are the putative *lin-4* homologs in mouse and human (9). miR-125a is detectable in PC-3, HeLa, Tera-2 (Figure 1), and other cell lines (data not shown). Two other miRNAs, miR-15 and miR-30, are present in a constant amount in these three cell lines (data not shown). In contrast, miR-125b level is highest in PC-3, moderate in HeLa and other cell lines, and undetectable in Tera-2 (Figure 1; data not shown). Tera-2 is an embryonic carcinoma (EC) cell line, representing undifferentiated cells. The higher expression of miR-125b in differentiated cells compared to undifferentiated cells is also seen with *let-7* (9,21,36-38). Tissues from prostate, cervix and brain have higher levels of these miRNAs than corresponding cell lines (Figure 1). Thus in all cell lines and tissues examined, human *let-7* and miR-125b are expressed specifically in differentiated cells or tissues, reminiscent of the higher expression seen later in development in flies and worms.

The depletion of miR-125b-2 in differentiated cancer cells inhibits cell proliferation

We first attempted to decrease miR-125b by siRNAs against the precursor of the mature miRNA. The hairpin shaped pre-miRNA is generated in the nucleus by the action of Drosha (39) but is exported to the cytoplasm (40) where it is processed by Dicer to release the mature miRNA (31,41-44). The pre-miRNA is therefore expected to be susceptible to siRNAs against the loop of the hairpin. There are two predicted precursors of miR-125b in humans, miR-125b-1 and -2. Both precursors share identical mature miRNA sequence, but the sequences flanking the mature miRNA are different (Figure 2A,

left panel). Two synthetic siRNAs were designed against the loop regions of miR-125b-1 and -2 (si125b-1 and si125b-2; Figure 2A, left panel). After transfection of these siRNAs into PC-3 cells, miR-125b level was measured by a primer extension assay. miR-125b was greatly reduced by the si125b-2 but not si125b-1 (Figure 2A, right panel). This observation suggests that most of the mature miR-125b is derived from the precursor miR-125b-2 in PC-3 cells.

To test if this strategy to knock-down miRNAs can be applied to other miRNAs, siRNAs against additional pre-miRNAs were transfected into PC-3 cells. miR-16 has two predicted precursors, miR-16-1 and -2. Transfection of siRNA against miR-16-1 and -2 (si16-1 and -2) decreased the mature miRNA to 40 % and 60 %, respectively (Figure 2B). Thus approximately 60 % of mature miR-16 is derived from the precursor -1 and the rest from precursor -2. Consistent with this interpretation, co-transfection of the two siRNAs against miR-16 decreased the level of miR-16 to approximately 20%, which is close to the background level. Efficient knock-down of miR-15a was also achieved by the transfection of siRNA against the loop region of the corresponding pre-miRNA.

Titration of siRNA dose revealed that siRNA concentration of 0.32 to 1 μ M was required to decrease the miRNA levels (Figure 2B). This dose is higher than the amount of siRNA required to knock-down a typical mRNA (0.02 to 0.3 μ M). The lower efficiency of the siRNA-directed degradation of a miRNA precursor could be due to the short length and/or transient nature of miRNA precursors. Indeed, the precursors of miRNA were never detected in our primer extension assays, presumably because miRNA precursors are quickly processed into mature miRNA. The inefficiency of siRNA against pre-miRNA might account for the failure of this strategy in two other cell lines, HeLa and

DU 145. Nevertheless, the results dictate that an siRNA can be used to deplete a miRNA in at least some cell lines like PC-3.

To investigate the biological effect of miR-125b, cell proliferation was measured after depletion of miR-125b by RNA interference (Figure 3A). si125b-2, but not si125b-1, reduced proliferation of PC-3 cells. Growth was restored by the co-introduction of a synthetic mature miR-125b that was designed to not anneal to si125b-2, demonstrating that growth suppression is specifically caused by the depletion of miR-125b. This result also indicates that miRNA function can be provided to cells by transient transfection of single stranded RNA. Tera-2 EC cells did not express miR-125b, and accordingly si125b-2 did not suppress growth of Tera-2 cells. To eliminate a trivial explanation, we demonstrated that the RNA interference machinery is intact in Tera-2 cells by transfecting siRNA against *ORC2* in this cell line and measuring Orc2 protein level (Figure 3B). Thus the failure of si125b-2 to suppress Tera-2 cell proliferation is not due to absence of the siRNA machinery and provides an additional control against non-specific effects of si125b-2 in PC-3 cells.

In order to confirm that the proliferation defect in Figure 3A is specifically caused by miR-125b depletion and to test if this defect is seen in cell lines other than PC-3, the experiment was repeated after transfection of 2'-O-methyl antisense oligonucleotide against miR-125b. Again, depletion of miR-125b by 2'-O-methyl oligonucleotide resulted in the reduction of proliferation, whereas the transfection of 2'-O-methyl oligonucleotide against GL2 did not (Figure 3C). In addition, this proliferation defect was observed not only in PC-3 cells, but also in HeLa cells. Since identical results were obtained from two independent methods and each experiment was performed with

stringent controls, it is clear that miR-125b is essential for the proliferation of differentiated cells.

miR-125b and *let-7* increase, while *lin-28* and *lin-41* decrease at mRNA level during differentiation

Taken together with the expression profiles of the human miRNAs in Figure 1, the previous results in *C. elegans* (17,19-23) suggest that *let-7* and miR-125b miRNAs are induced during differentiation and this upregulation of the miRNAs post-transcriptionally repress their putative targets which contain partially complementary target sequences in their 3' untranslated region. To test whether this regulation could be recapitulated in mammalian *in vitro* differentiation systems, we induced the differentiation of Tera-2 cells in the neuronal direction by retinoic acid (45). miR-125b became pronounced after 20 days of retinoic acid treatment (Figure 4, left panel). *let-7* increased to a level barely detectable after 20 days and to a higher level after 60 days. This time pattern was reminiscent of *C. elegans* in which *lin-4* is expressed earlier than *let-7* during development (17,19). In contrast to the miRNAs, Lin-28 protein decreased to barely detectable level after 20 days. Also, the mRNA of *lin-28* and *lin-41* disappeared completely by 20 days of retinoic acid (Figure 4, left panel). Thus the regulation of the miRNAs and their targets was recapitulated in the differentiation of EC cells *in vitro*.

To confirm the changes in expression of miRNAs and their targets in a second differentiation system, we induced the differentiation of mouse embryonic stem (ES) cells to the neuronal fate using the 5-stage *in vitro* differentiation procedure (34). Both

let-7 and miR-125b miRNAs were weakly expressed in ES cells, slightly decreased in embryoid bodies (EB), and then increased with differentiation through neuronal precursor (NP) stage into mature neurons (Figure 4, right panel). RNA from adult brain is shown for comparison. Lin-28 protein was expressed in ES and EB stages but absent in NP stage and later on. Notably, it was the mRNA level of *lin-28* and *lin-41* that is exactly anti-proportional to that of *let-7* and miR-125b (Figure 4, Northern blots in right panel). This suggests that miRNAs might regulate the mRNA level of *lin-28* and *lin-41* through posttranscriptional cleavage of mRNA as seen in plants, rather than regulate *lin-28* at the translational stage. It is worth noting that although there is a slight increase in *lin-28* mRNA and decrease in miR-125b miRNA between ES cells and EB, there was no increase in Lin-28 protein level in EB despite the decreased miRNA to target ratio. This, too, argues against translational control of *lin-28* by the miRNAs.

miR-125b and *let-7* are neither sufficient nor necessary for the transcriptional down-regulation of *lin-28* and *lin-41* during differentiation

Next, we attempted to provide a direct evidence for the effect of the miRNAs on the expression of their putative targets, by using our experimental tools to deplete or introduce a miRNA as described above. We performed a finer time course on the differentiating Tera-2 cells to test if there is an intermediate stage when the expression of *lin-28* and *lin-41* is regulated post-transcriptionally and is affected by experimental changes in miRNA levels. The steady-state mRNA levels of *lin-28* and *lin-41* started decreasing by 4-6 days, and were almost completely reduced by 12-14 days after retinoic acid treatment (Figure 5).

Since siRNA to the pre-miRNA did not work in the differentiating cells, we depleted miR-125b in this system by transfecting 2'-O-methyl antisense oligonucleotides (Figure 5A). In addition, RNA oligonucleotides with miR-125b and *let-7* sequence were transfected in order to assess the effect of premature induction of the miRNAs during differentiation (Figure 5B). Neither the depletion of miR-125b nor the premature introduction of mature miR-125b or *let-7* miRNA resulted in a striking change in the mRNA profile of *lin-28* and *lin-41* in the differentiating Tera-2 cells (Figure 5). Depletion of miR-125b produced a slight increase of *lin-28* mRNA at 8 and 14 days (Figure 5A, top panel lanes 9, 12), whereas transfection of miR-125b and *let-7* miRNA produced a slight decrease in *lin-28* mRNA (Figure 5B, top panel lanes 9, 10, 12, 13). Although these effects might suggest that the miRNAs repress the target transcriptionally or by post-transcriptional cleavage, the effects are marginal compared with the overall change of target message during differentiation.

Lin-28 protein remained relatively stable until 12-14 days of retinoic acid treatment (Figure 5A and B, bottom panel) followed by a significant reduction at 20 days and complete disappearance at 60 days (Figure 4, left panel). Thus the Lin-28 protein expression seems to simply follow the mRNA level during differentiation. Lin-28 protein level was also not affected by the depletion or introduction of miRNAs in the differentiating Tera-2 cells (Figure 5A and B, bottom panels). Transfection of *let-7* and miR-125b miRNAs did not decrease *lin-28* mRNA or protein in non-differentiating Tera-2 cells (data not shown). Conversely, depletion of miR-125b in PC-3 cells failed to induce *lin-28* mRNA or protein (data not shown). Translational regulation of Lin-41 by

let-7 is not plausible either, since *let-7* miRNA is barely detectable at 20 days by which time *lin-41* mRNA is fully down-regulated.

Taken together, these results suggest that at least in two mammalian *in vitro* differentiation models, down-regulation of *lin-41* and *lin-28* is not effected by translational suppression by *lin-4* and *let-7*. Although there is a decrease in Lin-28 protein during differentiation, the decrease correlates with a decrease in mRNA abundance of the *lin-28* gene.

***lin-28* and *lin-41*, specifically expressed in undifferentiated cells, are regulated at the transcriptional stage.**

Next, we investigated the expression of *lin-28* and *lin-41* in a number of human cell lines (Figure 6A) to generalize our result in Tera-2 differentiation. Among the cell lines tested, *lin-28* and *lin-41* mRNAs were observed only in Tera-2 which expressed no detectable level of *let-7* and miR-125b. All other cell lines have variable, but detectable level of *let-7* and miR-125b (data not shown) but with complete absence of *lin-28* and *lin-41* mRNA. Given the negative correlation in the expression of the miRNAs and their target mRNAs, it is not plausible that the miRNAs regulate the protein expression by annealing to the target mRNAs. Consistent with the abundance of *lin-28* and *lin-41* in undifferentiated cells, only testis expressed detectable levels of *lin-28* and *lin-41* mRNA among various tissues in a Northern blot (Figure 6B).

The low levels of *lin-28* and *lin-41* mRNAs could be due to decreased transcription or due to mRNA degradation directed by the miRNAs as seen in plants. Nuclear run-on assays were performed to address this question. Figure 6C and D show

that *lin-28* and *lin-41* are suppressed at the transcriptional level in all tested cell lines relative to the undifferentiated Tera-2 cells. Consistent with this, the decrease of *lin-41* mRNA on day 4 and of *lin-28* mRNA on day 8 in differentiating Tera-2 cells coincided with a reduction of transcription as measured by nuclear run-on assays (middle panel of Figure 5A). Therefore, the *in vitro* differentiation program appears to regulate the two genes mostly at the transcriptional level accounting for the unresponsiveness to any decrease or increase of miR125b or *let-7* miRNAs.

Discussion

siRNA to pre-miRNA to assess miRNA function.

With the discovery of hundreds of miRNAs in plants and animals methods that can quantitatively inactivate miRNAs are needed to assess their function. 2'-O-methyl oligoribonucleotides have been recently reported to specifically inactivate the RNAi activity of miRNAs in cultured cells and in *C. elegans* (26,27). Here we demonstrate that an siRNA against the loop region of the precursor to a miRNA can selectively downregulate the mature miRNA. The decrease of miRNA was monitored directly by primer extension assay whereas an inhibition of the miRNA can be monitored only through an indirect reporter-based assay when 2'-O-methyl antisense oligoribonucleotide is used.

The depletion of miR-125b specifically resulted in the proliferation defect and this proliferation defect was rescued by synthetic mature miRNA. Our result not only gives clues to the miRNA function but also provides an assay for miRNA function. Functional evaluation of a co-transfected mature miRNA is a particularly attractive aspect of this approach because such a rescue is hard to achieve when miRNAs are directly neutralized by 2'-O-methyl anti-sense oligonucleotides. This assay should allow us to evaluate sequence features of a miRNA that are critical for its function.

siRNA to miRNA precursor to assess miRNA biogenesis.

For many miRNAs computational methods have predicted multiple potential precursor genes in the genome. It is not clear whether all the precursor genes contribute equally to the mature miRNA pool. Since the loop regions of different pre-miRNAs are

not conserved, the use of siRNA against the loop region has a unique advantage over 2'-O-methyl antisense oligonucleotide in determining which miRNA precursor gene contributes to the mature miRNA pool in the cell. In the case of miR-16, both precursors contribute to the pool of mature miRNA in PC-3 cells. In contrast, miR-125b-2 is the primary source of the mature miR-125b in this cell, because si125b-2, but not si125b-1 depleted most of the mature miR-125b. The genes for miR-125b-1 and -2 are located in chromosome 11 and 21 respectively. miR-125b-2 is located in the intron of a gene while miR-125b-1 is located in the exon of a gene (NT_033899.473; Genscan). There was no significant difference in density of genes or of repetitive elements between these genomic loci. The Mfold program indicates that both precursors form hairpins with nearly identical stability. The difference in precursor use could be due to a difference in the transcription of the two genes containing the precursors. Alternatively a miRNA precursor in the intron of a gene may be more available for processing into a miRNA than a precursor contained in the exon of a gene. Further work is necessary to test these hypotheses, but the siRNA strategy allows us to begin addressing these questions.

Biological role of miR-125b in prostate cancer cell line, PC-3

The depletion of miR-125b through multiple transfections of si-125b2 resulted in a significant proliferation defect in PC-3 cells. The growth defect of PC-3 cells was not accompanied by an aberrant accumulation of cells in one stage of the cell cycle or by apoptosis (data not shown). The reduction of proliferation is not caused by a non-specific effect mediated by activation of PKR by the double-stranded siRNA (46), because it was not seen with si125b-1, nor was it seen in Tera-2 cells that did not express

the miRNA. Further, depletion of miR-125b by 2'-O-methyl oligonucleotide resulted in the same phenotype, supporting the specific effect of miR-125b on the cell proliferation. Several genes besides *lin-28* have been reported with mRNAs containing sequences that can form an RNA-RNA duplex with miR-125b (47,48). Possible involvement of these candidate genes is currently under investigation to account for the miR-125b knockout phenotype.

Conserved pattern of expression of miRNA targets, but diverged mechanism of regulation

We showed that *lin-28* and *lin-41* from human and mouse are specifically expressed in EC cells or ES cells in which miR-125b and *let-7* are not detectable. As differentiation proceeds, *lin-28* and *lin-41* decrease whereas miR-125b and *let-7* increase. This reciprocal expression pattern recapitulates the expression profile during *C. elegans* development. Our results are consistent with recently published reports suggesting that mammalian Lin-28 protein is down-regulated while the miRNAs miR-125b and *let-7* orthologs are up-regulated during differentiation (28,29). The conservation of sequence and expression pattern from nematode to humans suggests that the regulatory events mediated by *lin-4*, *let-7*, *lin-28*, and *lin-41* are also conserved.

There is however a marked divergence in the regulation of *lin-28* and *lin-41* between *C. elegans* and the mammalian *in vitro* differentiation systems studied here. In *C. elegans*, translational repression is the favored mechanism for the down-regulation of heterochronic genes by *let-7* and *lin-4*. As detailed in the introduction, a significant body of evidence leads us to expect a similar translational control of mammalian Lin-28 (28-

30). This expectation is particularly well supported by an experiment with an artificial luciferase construct fused to the *let-7* target sequence, naturally found in the 3' UTR of mammalian *lin-28* mRNAs (30). This construct is subject to *let-7* dependent translational repression strongly suggesting that the miRNAs can translationally repress the mammalian *lin-28* gene.

Our results, however, demonstrate that the regulation of mammalian *lin-28* and *lin-41* in the systems we studied is at the stage of transcription and is not affected by *let-7* and miR-125b. The nuclear run-on experiment during Tera-2 differentiation suggested that the down-regulation is transcriptional. In a panel of six cell lines the mammalian *lin-28* and *lin-41* appear regulated at the transcriptional stage. The mutual exclusivity in the expression of the miRNAs and their targets also supports the notion that the miRNAs do not regulate their targets at the translational stage. In addition, the artificial introduction or depletion of the miRNAs in differentiating or undifferentiated Tera-2 cells or PC-3 cells did not change the level of Lin-28 protein, *lin-28* mRNA, or *lin-41* mRNA, arguing against the possibility that the target mRNA is cleaved by the miRNA. This does not, of course, rule out the possibility of such mRNA cleavage in other situations. In our experiments, mouse ES cell was the only system in which the miRNAs and their targets exist together. However, here too, there was no increase in Lin-28 protein level in the subsequent EB stage relative to ES despite the decreased miRNA to *lin-28* mRNA ratio.

Thus siRNAs can downregulate miRNAs in mammalian cells in culture and we use them to (a) demonstrate an essential role of miR-125b in differentiated cells, (b) develop a functional assay for miR-125b and (c) identify which precursor gene is used to give rise to a miRNA. We also show that although the miRNAs miR-125b and *let-7* are

induced during differentiation of EC or ES cells in culture, while their targets *lin-28* and *lin-41* are repressed, the mechanism of the repression is different from the classical annealing of miRNA to target message and translation inhibition noted in *C. elegans*. *Lin-28* and *lin-41* are transcriptionally repressed and we have no evidence that the repression is mediated by the direct effect of miR-125b or *let-7*. Of course, these results do not imply that mammalian miR-125b and *let-7* never regulate their mammalian targets translationally. This question has to be re-visited if future experiments reveal additional examples (like the differentiating ES cells) where the target mRNAs and the miRNA are co-expressed, using the methods of miRNA depletion reported here.

In conclusion, we demonstrate that the miRNA depletion strategies used here are invaluable for discovering unexpected functions of miRNAs like the importance of miR125b in the viability of differentiated cells. In addition, such experiments are essential for objectively evaluating functions of miRNAs predicted from genetics in lower eukaryotes or from computational studies.

Acknowledgements

We would like to thank Dr. V. Narry Kim for helpful discussion, Dr. Eric Davies for kind advice on the ribosome profile experiment, and Dr. Deog Su Hwang for encouragement. This work was partly supported by a Post-doctoral Fellowship Program of Korea Science & Engineering Foundation (KOSEF) to YSL and a grant from the NIH (RO1 CA89406) to AD.

References

1. Ambros, V., Bartel, B., Bartel, D. P., Burge, C. B., Carrington, J. C., Chen, X., Dreyfuss, G., Eddy, S. R., Griffiths-Jones, S., Marshall, M., Matzke, M., Ruvkun, G., and Tuschl, T. (2003) *Rna* **9**, 277-279
2. Lee, R. C., Feinbaum, R. L., and Ambros, V. (1993) *Cell* **75**, 843-854
3. Lee, R. C., and Ambros, V. (2001) *Science* **294**, 862-864
4. Lagos-Quintana, M., Rauhut, R., Lendeckel, W., and Tuschl, T. (2001) *Science* **294**, 853-858
5. Lau, N. C., Lim, L. P., Weinstein, E. G., and Bartel, D. P. (2001) *Science* **294**, 858-862
6. Dostie, J., Mourelatos, Z., Yang, M., Sharma, A., and Dreyfuss, G. (2003) *Rna* **9**, 180-186
7. Grad, Y., Aach, J., Hayes, G. D., Reinhart, B. J., Church, G. M., Ruvkun, G., and Kim, J. (2003) *Mol Cell* **11**, 1253-1263
8. Lagos-Quintana, M., Rauhut, R., Meyer, J., Borkhardt, A., and Tuschl, T. (2003) *Rna* **9**, 175-179
9. Lagos-Quintana, M., Rauhut, R., Yalcin, A., Meyer, J., Lendeckel, W., and Tuschl, T. (2002) *Curr Biol* **12**, 735-739
10. Lim, L. P., Glasner, M. E., Yekta, S., Burge, C. B., and Bartel, D. P. (2003) *Science* **299**, 1540
11. Mourelatos, Z., Dostie, J., Paushkin, S., Sharma, A., Charroux, B., Abel, L., Rappsilber, J., Mann, M., and Dreyfuss, G. (2002) *Genes Dev* **16**, 720-728
12. Park, W., Li, J., Song, R., Messing, J., and Chen, X. (2002) *Curr Biol* **12**, 1484-1495
13. Reinhart, B. J., Weinstein, E. G., Rhoades, M. W., Bartel, B., and Bartel, D. P. (2002) *Genes Dev* **16**, 1616-1626
14. Llave, C., Kasschau, K. D., Rector, M. A., and Carrington, J. C. (2002) *Plant Cell* **14**, 1605-1619
15. Ambros, V., Lee, R. C., Lavanway, A., Williams, P. T., and Jewell, D. (2003) *Curr Biol* **13**, 807-818
16. Suh, M. R., Lee, Y., Kim, J. Y., Kim, S. K., Moon, S. H., Lee, J. Y., Cha, K. Y., Chung, H. M., Yoon, H. S., Moon, S. Y., Kim, V. N., and Kim, K. S. (2004) *Dev Biol* **270**, 488-498
17. Reinhart, B. J., Slack, F. J., Basson, M., Pasquinelli, A. E., Bettinger, J. C., Rougvie, A. E., Horvitz, H. R., and Ruvkun, G. (2000) *Nature* **403**, 901-906
18. Pasquinelli, A. E., and Ruvkun, G. (2002) *Annu Rev Cell Dev Biol* **18**, 495-513
19. Feinbaum, R., and Ambros, V. (1999) *Dev Biol* **210**, 87-95
20. Moss, E. G., Lee, R. C., and Ambros, V. (1997) *Cell* **88**, 637-646
21. Pasquinelli, A. E., Reinhart, B. J., Slack, F., Martindale, M. Q., Kuroda, M. I., Maller, B., Hayward, D. C., Ball, E. E., Degan, B., Muller, P., Spring, J., Srinivasan, A., Fishman, M., Finnerty, J., Corbo, J., Levine, M., Leahy, P., Davidson, E., and Ruvkun, G. (2000) *Nature* **408**, 86-89
22. Slack, F. J., Basson, M., Liu, Z., Ambros, V., Horvitz, H. R., and Ruvkun, G. (2000) *Mol Cell* **5**, 659-669

23. Vella, M. C., Choi, E.-Y., Lin, S.-Y., Reinert, K., and Slack, F. J. (2004) *Genes Dev.* **18**, 132-137
24. Kawasaki, H., and Taira, K. (2003) *Nature* **423**, 838-842
25. Kawasaki, H., and Taira, K. (2003) *Nature* **426**, 100
26. Meister, G., Landthaler, M., Dorsett, Y., and Tuschl, T. (2004) *Rna* **10**, 544-550
27. Hutvagner, G., Simard, M. J., Mello, C. C., and Zamore, P. D. (2004) *PLoS Biol* **2**, E98
28. Moss, E. G., and Tang, L. (2003) *Developmental Biology* **258**, 432-442
29. Sempere, L. F., Freemantle, S., Pitha-Rowe, I., Moss, E., Dmitrovsky, E., and Ambros, V. (2004) *Genome Biol* **5**, R13
30. Nelson, P. T., Hatzigeorgiou, A. G., and Mourelatos, Z. (2004) *Rna* **10**, 387-394
31. Hutvagner, G., McLachlan, J., Pasquinelli, A. E., Balint, E., Tuschl, T., and Zamore, P. D. (2001) *Science* **293**, 834-838
32. Sambrook, J., Fritsch, E. F., and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY
33. Pertovaara, L., Tienari, J., Vainikka, S., Partanen, J., Saksela, O., Lehtonen, E., and Alitalo, K. (1993) *Biochem Biophys Res Commun* **191**, 149-156
34. Lee, S. H., Lumelsky, N., Studer, L., Auerbach, J. M., and McKay, R. D. (2000) *Nat Biotechnol* **18**, 675-679
35. Greenberg, M. E., and Bender, T. P. (2002) in *Short Protocols in Molecular Biology* (Struhl, K., ed) Vol. 1, pp. 4-25 - 24-29, 2 vols., John Wiley & Sons, Inc.
36. Sempere, L. F., Sokol, N. S., Dubrovsky, E. B., Berger, E. M., and Ambros, V. (2003) *Dev Biol* **259**, 9-18
37. Bashirullah, A., Pasquinelli, A. E., Kiger, A. A., Perrimon, N., Ruvkun, G., and Thummel, C. S. (2003) *Dev Biol* **259**, 1-8
38. Aravin, A. A., Lagos-Quintana, M., Yalcin, A., Zavolan, M., Marks, D., Snyder, B., Gaasterland, T., Meyer, J., and Tuschl, T. (2003) *Dev Cell* **5**, 337-350
39. Lee, Y., Ahn, C., Han, J., Choi, H., Kim, J., Yim, J., Lee, J., Provost, P., Radmark, O., Kim, S., and Kim, V. N. (2003) *Nature* **425**, 415-419
40. Lund, E., Guttinger, S., Calado, A., Dahlberg, J. E., and Kutay, U. (2004) *Science* **303**, 95-98
41. Ketting, R. F., Fischer, S. E., Bernstein, E., Sijen, T., Hannon, G. J., and Plasterk, R. H. (2001) *Genes Dev* **15**, 2654-2659
42. Bernstein, E., Caudy, A. A., Hammond, S. M., and Hannon, G. J. (2001) *Nature* **409**, 363-366
43. Grishok, A., Pasquinelli, A. E., Conte, D., Li, N., Parrish, S., Ha, I., Baillie, D. L., Fire, A., Ruvkun, G., and Mello, C. C. (2001) *Cell* **106**, 23-34
44. Knight, S. W., and Bass, B. L. (2001) *Science* **293**, 2269-2271
45. Thompson, S., Stern, P. L., Webb, M., Walsh, F. S., Engstrom, W., Evans, E. P., Shi, W. K., Hopkins, B., and Graham, C. F. (1984) *J Cell Sci* **72**, 37-64
46. Balachandran, S., Kim, C. N., Yeh, W. C., Mak, T. W., Bhalla, K., and Barber, G. N. (1998) *Embo J* **17**, 6888-6902
47. Lewis, B. P., Shih, I. H., Jones-Rhoades, M. W., Bartel, D. P., and Burge, C. B. (2003) *Cell* **115**, 787-798

48. John, B., Enright, A. J., Aravin, A., Tuschl, T., Sander, C., and Marks, D. S. (2004) *PLoS Biol* **2**, e363

Figure Legends

Figure 1. Quantitative measurement of miRNAs from various cell lines and tissues

Primer extension assays were performed with RNAs from various cell lines and tissues as described in Materials and Methods, to detect *let-7*, miR-125a, miR-125b, and 5S rRNA as a control. The extension products are indicated by arrows.

Figure 2. Depletion of miR-125b and two other miRNAs by siRNAs against the precursors

A. The two hairpin precursors from predicted for miR-125b are shown in the left panel. Underlined black: mature miR-125b sequence. Capitals black: the target sequences of siRNAs. Right panel shows the primer extension assays to detect miR-125b or 5S rRNA control after three rounds of transfection at 24 hours intervals with each siRNA.

B. Quantitation of miRNAs (right) by primer extension after transfection of the siRNAs (left). The concentration of each siRNA during the transfection is indicated in parenthesis. The relative intensity of the miRNA primer extension-product was normalized to that of the 5S rRNA control and compared to the corresponding signal from mock-transfected cells.

Figure 3. The effect of miR-125b depletion on the proliferation of differentiated cells.

A. After transfection of miR-125b and/or siRNA against miR-125b into PC-3 and Tera-2 cells, cell proliferation was measured by MTT assay. First three transfections were done

at 24 hours intervals, fourth transfection onwards were at 48 hours intervals. X-axis indicates the number of transfections, Y-axis OD value at 570 nm.

B. Immunoblot of Orc2 protein (indicated by arrow) from Tera-2 cells transfected without (mock) or with siRNA against *orc2* (si-*orc2*)

C. After transfection of 2'-O-methyl antisense oligonucleotide against miR-125b and GL2 control, cell proliferation of PC-3 and HeLa was measured as described in panel A.

Figure 4. Expression of miR-125b, *let-7*, and their targets during differentiation of cells in culture.

left panel; Primer extension assay (*let-7*, miR-125b), Northern hybridization (*lin-41*, *lin-28*, *gapdh*) and immunoblot (Lin-28, β -actin) were performed with samples prepared from Tera-2 embryonic carcinoma cells before (0 days) or after 20 and 60 days of retinoic acid (RA) treatment. *right panel*; Same assays were performed with samples from mouse ES cells subjected to *in vitro* differentiation. ES, embryonic stem cells; EB, embryoid body; NP, neuronal precursor; 5:10 and 5:15, 10 and 15 days of the final neuronal differentiation stage (stage 5). In Lin-28 immunoblot, equal loading of protein in each lane was confirmed by Ponceau S staining of the blot (not shown). Total RNA from brain was included as a positive control for differentiated tissue.

Figure 5. Effect of miRNAs on putative targets during *in vitro* differentiation

A. Northern hybridization (top panel), nuclear run-on (middle) and immunoblot (bottom) were performed with samples prepared from Tera-2 cells after 4 days without retinoic acid (RA) treatment (w/o) or after 4, 8, and 14 days of retinoic acid (RA) treatment. The

cells were not transfected (-) or transfected with 2'-O-methyl antisense oligonucleotide against GL2 (GL2) and miR-125b (125b). First three transfections were done at 24 hours intervals and fourth transfection onwards were at 48 hours intervals. Nuclear run-on experiment was performed at the same time points during differentiation with untransfected Tera-2 cells. In Lin-28 immunoblot, Ponceau S staining of the blot shows equal loading of protein in each lane.

B. Northern hybridization (top panel) and immunoblot (bottom) were performed with samples prepared before (0 days) or after 3, 6, 9, and 12 days of retinoic acid (RA) treatment combined with the transfections of RNA oligonucleotides encoding miR-125b or *let-7*.

Figure 6. Expression of *lin-28* and *lin-41*, putative targets of the miRNAs, in various cell lines and tissues.

A. mRNA levels of *lin-28* and *lin-41*, as well as *gapdh* (glyceraldehyde-3-phosphate dehydrogenase) as a control, in various cell lines were detected by Northern hybridizations. The locations of 18S and 28S ribosomal RNA are indicated in the left.

B. Northern hybridizations were performed on a tissue blot (human MTN blot II from Clontech).

C. Nuclear run-on experiments were performed with nuclei from the indicated cell lines. The identities of each band are marked in a box below. pBS indicates the pBluescript vector negative control. 293T cells have an integrated pBluescript.

D. Each band in the blot of panel E was quantitated by autoradiography. Signal from pBluescript vector negative control was subtracted and then the intensities were

normalized to that for the *gapdh* gene. In the case of 293T, signal from the blank area was used as background while normalization instead of that from pBluescript vector.

Figure 1




Figure 2B




Figure 3A





Figure 3B




Figure 3C




Figure 4




Figure 5A




Figure 5B




Figure 6A




Figure 6B




Figure 6C




Figure 6D

